

1
2
3
4 **COPING WITH CLIMATE CHANGE: THE ROLE OF**
5
6 **SPATIAL DECISION SUPPORT TOOLS IN FACILITATING**
7
8 **COMMUNITY ADAPTATION**
9

10
11
12 DAVID J. LIESKE^{1,2}
13
14

15
16
17 ¹Department of Geography and Environment, Mount Allison University, 144 Main Street,
18 Sackville, New Brunswick, Canada
19

20
21
22 ²Corresponding author: dlieske@mta.ca
23
24
25

26 **Abstract**
27

28
29 Climate change challenges communities to visualize spatial patterns of risk, assess their
30 vulnerability to those risks, and prepare adaptation plans to lower vulnerability. This paper
31 outlines the design and implementation of a prototype web-based spatial decision support
32 system (SDSS), referred to as the Community Adaptation Viewer (CAV), to assist adaptation
33 planning. Thin-client, Javascript enabled web-SDSS software was constructed to allow
34 interaction with urban infrastructure, and support "on-the-fly" assessment of social and
35 economic vulnerability. Facilitated, decision-making workshops were conducted with small
36 groups of stakeholders to evaluate the effectiveness of the prototype. The test case illustrates
37 that high levels of information integration are practical to achieve, and that SDSS can
38 significantly enhance the ability of communities to conduct elaborate, geographically-specific
39 climate change adaptation planning. Given the long time frame required to fulfill some
40 adaptation plans, it is crucial that communities begin to develop and invest in adaptation
41 strategies as soon as possible.
42
43
44
45
46
47
48
49
50
51

52
53
54
55
56 **Keywords:** web-based software, spatial decision support, climate change, adaptation,
57 flood risk
58
59
60
61
62
63
64
65

Software Availability

Name of Software:	Community Adaptation Viewer (CAV)
Developer:	David J. Lieske
Contact Address:	Dept. of Geography and Environment Mount Allison University 144 Main Street, Sackville, New Brunswick Canada, E4L 1A7
Contact E-mail:	dlieske@mta.ca
Available Since:	2013
Programming Language:	Javascript
Availability:	http://arccgis.mta.ca/toolkit
Cost:	Free

1. Introduction

Decision analysis centers on taking problems, dividing them into small, more manageable parts, analyzing each of those parts, and integrating the findings to produce logically meaningful solutions (Karnatak et al. 2007). In the case of classic geospatial problems such as network routing, the goal consists of taking a starting point and identifying the least cost-path (or shortest commute time) to a set of destinations given a particular road network. The problem lends itself to complete parameterization, and yields a finite and very clearly defined optimal solution (or set of solutions).

Planning for climate change adaptation falls outside of this type of well-structured problem domain. In fact, with the implicit aim of reducing community vulnerability, climate change adaptation is "ill structured" (Sugumaran and DeGroot 2011) in a number of ways. First, it involves complex social, environmental and economic dimensions which are not easy to quantify. In the case of social vulnerability, there is a wide consensus that it is, in part, a socially-constructed consequence of uneven access to economic, social, and informational resources (Cutter et al. 2000, Wu et al. 2002, National Research Council 2006, Füssel 2007,

1
2
3
4 Hebb and Mortsch 2007, Wilby and Keenan 2012). Assuming that a set of measurable social
5 vulnerability indicators are agreed upon to capture this reality, there are intangibles presented
6 by the socio-political landscape that will impinge upon and determine the acceptability of any
7 potential risk reduction solution. Other facets of the climate change adaptation problem
8 include the fact that the goals and objectives of community stakeholders may not be
9 completely definable, even in the minds of the stakeholders themselves. Furthermore, the
10 goals may be competing or even in opposition to each other. We can also expect there to be
11 many possible candidate solutions, the efficacy of which will be plagued by uncertainty.
12
13

14
15 In their analysis of adaptation planning in agricultural contexts, Risbey et al. (1999)
16 identified a number of aspects of the decision-making process that have wide utility for
17 understanding the framework in which adaptation decision making occurs. *Signal detection*,
18 which involves perception of relevant patterns (e.g., trends in time or space, Chen 2005) is a
19 necessary prerequisite. SDSS and visualization tools have an obvious role to play here
20 (Eppler and Aeschimann 2009), allowing the mapping of risk and information, and detection
21 of temporal trends in environmental conditions. Models can be used to depict the outcome
22 of different climate change scenarios, and animations can provide powerful impressions of
23 particular events. For instance, infrastructure impacts can be depicted using computer-
24 generated graphics or historical images, which studies have shown can strongly impact risk
25 perception and deepen understanding of the problem (Lieske et al. 2014).
26
27

28
29 The second aspect of adaptation planning identified by Risbey et al. (1999) is *evaluation*,
30 where the interpretation of future projections and assessment of the foreseeable impacts are
31 conducted. In climate change adaptation, this constitutes "vulnerability analysis", and
32 involves the identification of at-risk "hot spots" along with the economic, social and
33 environmental implications. In addition to deepening understanding of vulnerabilities, this
34 analysis is a necessary prerequisite to prioritization of the right courses of action.
35
36

37
38 The final phases of Risbey et al. (1999) involve *decision and response*, and *feedback*, both of
39 which could be greatly augmented by SDSS. In addition to allowing for the simultaneous
40 assessment of the potential lowering of vulnerability with respect to implementation cost,
41 SDSS can act as a living database of the risk appraisal process, inviting re-analysis and
42 reflection at any point. Different audiences or constituencies can use the SDSS to draw their
43 own conclusions, thereby diversifying the portfolio of possible risk reduction strategies as
44 well as flagging different aspects of community vulnerability. The creation of user-identified
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 features and annotations within SDSS software acts as a record ("shared memory") inviting
5 further and subsequent analysis (Andrienko et al. 2007), thereby preserving the 'derived
6 knowledge" of the adaptation planning process (Peuquet and Kraak 2002, Mennis and
7 Peuquet 2003, Andrienko et al. 2007, Hopfer and MacEachren 2007).
8
9

10
11 In order for communities to make meaningful progress in planning for and
12 implementing necessary short- and long-term changes, there is a need for information
13 systems which allow community stakeholders to visualize climate change risk information
14 together with various aspects of community vulnerability (Flax et al. 2002). Such systems
15 should work in group settings and encourage collaboration by diverse groups of stakeholders
16 of differing competencies, interests, and political agendas (Jankowski et al. 1997); yet, they
17 should also help users to discern critically vulnerable locations, better understand the risks
18 and vulnerabilities involved, and create plausible scenarios representing multiple courses of
19 action. By assisting in the computation and visualization of the impacts of various adaptation
20 measures, it should help users to articulate goals and priorities, thereby identifying the right
21 course of action for the community. Software has tremendous potential to provide the
22 information framework necessary to support all aspects of adaptation planning, and to
23 "complement the power of computational methods with human background knowledge,
24 flexible thinking, imagination, and capacity for insight" (Andrienko 2007: 840). For reasons I
25 argue below, a new generation of spatial decision support system (SDSS) tools have the
26 potential to deliver this support.
27
28
29
30
31
32
33
34
35
36
37
38
39

40 This paper reports on a project that originated with the findings of earlier climate-
41 change visualization work, in which a need for software capable of displaying risk and
42 sensitivity layers, along with the interactive identification of locations of concern (LOCs) and
43 candidate adaptation planning zones (APZs), was identified early on. In order to maximize
44 the accessibility and ease-of-use of the software, a design decision was made to implement a
45 prototype web-based spatial decision support system (web-SDSS) using a thin-client (see
46 Power and Kaparathi 2002), Javascript user interface. This conferred significant advantages,
47 such as the central storage of data and ability to run in ordinary web browsers. The software,
48 referred to as the "Community Adaptation Viewer" (CAV), also provided functions for
49 assessing economic and social vulnerability "on-the-fly" at a high level of detail.
50
51
52
53
54
55
56
57

58 The second phase of this project involved test deployment of the software in a
59 community in south-east New Brunswick, Canada, which is at significant risk to sea-level
60
61
62
63
64
65

1
2
3
4 rise-related coastal flooding. Through the use of facilitated workshops, small groups of
5 stakeholders were brought together to use the CAV. This allowed a progression from
6 problem identification and characterization to adaptation planning zone delineation, and
7 resulted in the brainstorming of possible risk reduction strategies. This paper reviews both
8 the software and the community case study, and offers suggestions for how this (or similar)
9 software can help advance the world-wide adaptation agenda.
10
11
12
13
14
15
16
17

18 **2. Design Considerations**

19
20
21

22 Due to time and manpower constraints, as well as limitations in the availability of
23 project stakeholders, development could only be described as partially user-centred (see
24 Maguire 2001, Gulliksen et al. 2003). Users were not involved in all stages of project
25 development (as would be expected in a true user-centred iterative approach), but data
26 requirements and overall functionality was derived from feedback received during two earlier
27 phases of consultation: (1) seven meetings held between November 5 and 18, 2013 to
28 identify and discuss the concept of "vulnerability", and (2) an earlier study involving ten
29 attendees from six of the twelve then-existing provincial planning districts (Rones 2013).
30 The flexibility of web-based software was uniformly appreciated, and functionality such as
31 ability to estimate the economic costs of potential flood events was suggested as key
32 information to support decision making. Previous efforts excluded social vulnerability
33 indicators, so this was considered an important data component that needed to be
34 developed.
35
36
37
38
39
40
41
42
43
44

45 A design decision was made to adopt a "thin client" approach (see Power and Kaparthy
46 2002, Rinner 2003) and build a web-enabled user interface using the Javascript API and
47 Dojo Toolkit. As pointed out by Chen et al. (2007), this eliminates the need for users to have
48 to download and install software or manage updates ("thick client"), thereby increasing
49 flexibility and long-term usability (Bhargava et al. 2007). Previous work has argued that
50 interactivity enhances the quality of generated solutions (Andrienko and Andrienko 2006,
51 Andrienko et al. 2007) so a high level of user-interface interactivity was considered necessary.
52 Interface development proceeded in accordance with Schneiderman's "Information Seeking
53 Mantra" of *overview first, zoom and filter, and then details-on-demand* (Schneiderman 1996), allowing
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 users to actively guide/influence the analysis, reduce the breadth and/or depth of the search
5 (saving on computation time) and focus attention on the relevant output.
6

7
8 Real-time responsiveness was considered an important design goal given the need for
9 users to be able to receive immediate feedback in an "appropriate" form (Andrienko et al.
10 2007). In order to work through the implications of the emerging community vulnerability
11 patterns (the "hot spots") during the course of a typical focus group, users must be able to
12 rapidly assess the risk exposure in these areas.
13
14

15
16 Underlying spatial data (e.g., lidar used to accurately measure surface elevation at 1-m
17 spatial resolution and produce maps of potential flood extents) were housed in a single,
18 integrated geodatabase to be hosted as map service layers on ArcGIS Server 10.1 (ESRI
19 2014). Infrastructure at-risk (e.g., highways, institutions, properties) were also gathered
20 within the same database, and included other information such as potential economic
21 impacts and social vulnerability information. This approach centralized important risk and
22 vulnerability layers, and rendered them accessible from any location connected to the
23 internet.
24
25

26
27 Finally, visualization research has pointed to the importance of storing the valuable,
28 intangible, and largely unanticipated knowledge provided by users (Peuquet and Kraak 2002,
29 Andrienko et al. 2007). Referred to as "shared memory" (Andrienko et al. 2007) or "derived
30 knowledge" (Mennis and Peuquet 2003), web-accessible annotation tools are considered an
31 ideal way to gather such data (Hopfer and MacEachren 2007). The need to support user-
32 provided data is expected to be especially crucial in climate change vulnerability assessments
33 when expert knowledge is available to supplement existing information. For spatial decision
34 support systems, this requires tools for drawing point and areal features, assigning attribute
35 information to those features, and storing them in the centralized database.
36
37

38
39 As a final design note: while ultimately not pursued due to time and resource
40 limitations, previous research has pointed to the value of archiving hyperlinked information,
41 e.g., pictures of affected or vulnerable areas (Aggett and McColl 2006), climate change
42 projections, IPCC reports, and model simulations/animations. This ensures that supporting
43 documentation and resources are centrally gathered and accessible, and can potentially
44 extend the utility of climate change adaptation software. However, developers need to
45 balance the temptation to extend the functionality of SDSS tools with the need to efficiently
46 address key project goals.
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

3. System Architecture

The CAV architecture consists of three interlocking layers: a data layer (consisting of physical infrastructure as well as economic and social vulnerability data), a visualization layer (the GUI interface), and a processing layer (Fig. 1).




Figure 1. System architecture of the Community Adaptation Viewer (CAV).

1
2
3
4
5
6 3.1. Data Layer
7
8
9

10 Physical infrastructure data (e.g., roads, schools, high density housing; Table 1) were
11 compiled to support potential user-directed queries. Some information was obtained directly
12 from the Province of New Brunswick's geospatial data gateway
13 (<http://www.snb.ca/geonb1/e/index-E.asp>), while others were derived products (e.g.,
14 building footprints created by digitizing orthophotos).
15
16
17
18
19

20 Table 1.

21
22 **Anticipated Information Requirements**

23

24 Layer Group	25 Data	26 Resolution / 27 Scale	28 Source	29 Description
30 Dyke 31 Vulnerability	32 Height	33 10 m	34 MTA GML ^a	35 Maximum 36 height and 37 elevation 38 profile at each 39 10 m section 40 of dyke. 41 Derived from 42 LiDAR.
	43 Susceptibility 44 to erosion	45 10 m	46 MTA GML	47 Modelled 48 from NDVI, 49 proximity to 50 water
51 Social 52 Vulnerability	53 Social 54 Vulnerability 55 Index (SVI)	56 Dissemination 57 Area (area 58 containing 59 ~500 persons)	60 http://www.statcan.gc.ca/	61 Social 62 vulnerability 63 obtained using 64 Stats Canada 65 Census 66 Dissemination 67 Areas
68 Infrastructure 69 Vulnerability	70 Roads	71 1:10,000	72 RSC7 ^b planning	73 Digitized 74 from 2013 75 satellite 76 imagery and 77 SNB DPM
	78 Hospitals	79 0.05-10 m	80 SNB ^c real property 81 attribute data	

82
83
84
85

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

	Seniors residences	0.05-10 m		
	High density residences	0.05-10 m	SNB	
	Building Footprints	1:5000	MTA GML	Digitized from 2009 orthophotos
	Historic sites		Parks Canada	
	Waste treatment facility	0.05-10 m	SNB	
	Lift Station	0.05-10 m	SNB	
	School		SNB	
	Church	0.05-10 m	SNB	
	Storm water system	0.5-10 m	Town of Sackville/MTA GML	
	Rail Line	0.05-10 m	SNB	
Economic Vulnerability	Tax Assessment Values (Province of NB)	0.05-10 m	SNB	Privacy concerns require that individual properties can- not be displayed
Land Allocation	AgUse	1:10,000?	NB Dept. Aquaculture Agriculture and Fisheries	
	Zoning	0.05-10 m	RSC7 planning	
Adaptation Planning	Point of Concern (POC) Adaptation Planning Zone (APZ)			User-created during workshops User-created during workshops

^aMount Allison University, Geospatial Modelling Lab.
^bRegional Service Commission 7 (<http://www.nbse.ca/home>).
^cService New Brunswick, geographic information gateway (<http://www.snb.ca/geonb1/e/index-E.asp>).

Map layers are served using the ArcGIS Server REST API (ESRI 2014), which provides a simple and open (though stateless) web interface to map services through a

1
2
3
4 hierarchy of uniform resource locators (URLs). Other services supported by the REST API
5 include geoprocessing services, which perform server-side geoprocessing tasks, and
6 geometry and feature services, for drawing and interacting with geodatabase feature objects.
7
8 Feature objects used in this project included locations of concern (LOC) and adaptation
9 planning zones (APZs), discussed in Section 3.2.
10
11

12
13 The integration of economic (Section 3.1.1.) and social vulnerability information
14 (Section 3.1.2) was an important design criteria for the CAV, as such information is normally
15 scattered or unavailable. However, special processing was required to prepare this data.
16
17
18
19

20 *3.1.1. Quantification of Economic Vulnerability*

21
22
23

24 Flood damage information, based on the work of Wilson et al. (2012), was calculated
25 for current (2000) and projected (2025, 2055, 2085 and 2100) 1-in-100 year sea levels.
26 Damage depth curves relate the depth of floodwater to the expected severity of damage, and
27 were obtained from the U.S. Army Corps of Engineers. The damage costs are based on
28 flood depth, property value, and are weighted by the percentage of the building footprint
29 flooded on any given parcel. Vehicle costs were only calculated for residential buildings that
30 were flooded and number of vehicles was determined from Natural Resources Canada
31 (NRCAN) assessment of NB households, and market values assessed from a review of local
32 used vehicle prices. Total economic vulnerability was estimated by summing the value of
33 exposed building structures, building contents, vehicle values, and (where relevant)
34 agricultural crop values. Agricultural damages considered the value of the crop type at
35 different times of year, and the percentage of active agricultural land flooded. It was assumed
36 that damages to residential, commercial, and public parcels are tied to the structures on the
37 parcels.
38
39
40
41
42
43
44
45
46
47
48

49 It should be noted that estimates of repair costs to public infrastructure (e.g., roads,
50 bridges, culverts) were unavailable. For this reason, economic vulnerability should be treated
51 as an underestimate of the likely damage costs.
52
53
54
55

56 *3.1.2. Quantification of Social Vulnerability*

57
58
59
60
61
62
63
64
65

1
2
3
4 Social vulnerability was assessed using a social vulnerability index (SVI) to reflect
5 residents' relative ability to prepare, respond and recover from flooding. When creating the
6 SVI, care was taken to ensure that the range of social vulnerabilities to flooding were
7 reflected, that the vulnerability types were equally weighted, and that the index was easily
8 interpretable.
9

10
11
12
13 King and MacGregor (2000), Cutter et al. (2000) and Wu et al. (2002) identified the
14 types of social vulnerabilities that can make it difficult for neighbourhoods to prepare,
15 respond and recover from flooding. Analogues of these indicators were obtained from the
16 2006 Canada Census (Statistics Canada), and Principal Component Analysis (PCA) was used
17 to group similar, or correlated, indicators together. Interpretation of the PCA shows that
18 four socially-vulnerability groups exist in New Brunswick: socioeconomic status (wealth,
19 knowledge and employment status), minorities and renters, elderly, and youth.
20

21
22
23
24
25
26 When creating the SVI, a single representative measure was selected from each of the 4
27 vulnerability groups: median income, percentage of dwellings that are rented, aged
28 dependency (the ratio of persons over 65 to the population aged 15 to 65) and youth
29 dependency (the ratio of persons under 15 to the population aged 15 to 65). Renters were
30 selected instead of minorities because in the Town of Sackville renters were deemed to be
31 more vulnerable. Each vulnerability measure was scaled out of 1 (by dividing by the
32 provincial maximum value), and the sum of the 4 scaled measures was calculated to create
33 the SVI. A SVI score of near to 0 means that communities are less vulnerable to flooding
34 while a SVI score approaching 4 are more vulnerable.
35
36
37
38
39
40
41
42
43
44


45 *3.2. Visualization Layer*

46
47
48

49 The visualization layer (Fig. 1) consists of a GUI developed using the Javascript API
50 (Flanagan 2011). The Javascript API is a good choice for developing client-side interfaces
51 given that it is a language deployed throughout the world wide web and highly integrated
52 with web browser objects (Power and Kaparathi 2002). This relieves users of the burden of
53 having to install and update custom software on their local workstations (Sugumaran and
54 DeGroot 2011). Projects such as the Dojo toolkit (www.dojotoolkit.org, Russell 2008)
55 extend the functionality and utility of Javascript, allowing the production of special graphical
56
57
58
59
60
61
62
63
64
65

1
2
3
4 output, for instance. Dojo is used extensively by ESRI's Javascript API to govern interaction
5 with the ArcGIS Server.
6

7
8 The main map viewer pane (Fig. 2) displays both a basemap, as well as any and all data
9 layers activated by the user using check box controls housed within the "Layer Selection"
10 of the accordian pane (Fig. 2). The "Basemaps" button on the task pane (Fig. 2) calls the
11 BasemapGallery customized Dojo dijit (ESRI 2014), allowing the selection of nine
12 different basemap layers. Subsequent user feedback indicated that the use of an aerial
13 imagery baselayer, while displaying land cover details, could be disorienting. Pan and zoom
14 functionality is automatically built into the Map object class, and triggered by click and
15 mouse wheel events.
16
17
18
19
20
21
22



23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52 **Figure 2.** Schema of the main user interface.

53
54
55
56 Feature creation is launched by clicking on the appropriate icon in the feature-creation
57 pane (Fig. 2). Users have a choice of two types of geometry: a point geometry feature
58 (locations of concern, LOC; Fig. 3a) or two classes of areal geometry (adaptation planning
59
60
61
62
63
64
65

1
2
3
4 zones, APZ; dyke improvement zone, DIZ; Fig. 3b). At any time users can access and edit
5 the attribute information pertaining to that feature, or delete features altogether (Fig. 3). The
6 SessionID field stores a session-specific identifier for that feature, ensuring that only
7 features created during a single adaptation planning session are displayed in the map viewer
8 pane. Data from other sessions are hidden, unless "all" is specified at the time of page
9 loading.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 3. Text controls for editing attribute information for user-created locations of concern (a), and adaptation planning zones (b).

1
2
3
4 Project metadata is exposed via the "Further Information" tab of the accordion pane,
5 which presents a pulldown control with a fixed set of optional topics. XMLHttpRequest
6 function calls are used to open simple HTML text files hosted on the Documentation Server
7 (Fig. 1). For the CAV project, topics included information on sea-level projections, as well as
8 the social vulnerability index.
9

10
11
12
13 Once users have explored the map viewer and created APZ features, the "Select
14 Planzones" button on the task pane (Fig. 2) permits the selection of one or more APZ
15 features using an expandable window frame. "Selection" status is indicated using a broken
16 red border and yellow shading, which corresponds to the selected geodatabase features
17 contained within the Database server (Fig. 1). Selected features can be deselected using the
18 "Clear Selection" button on the task pane (Fig. 2), or passed on to the Processing Layer (Fig.
19 1) by clicking on the "Analyze!" button (Fig. 2). Example of geoprocessing output is shown
20 in Figure 4. Figure 4 (a) is a barchart widget summarizing total estimated economic
21 vulnerability for all parcels intersecting the selected APZ features, under each of five
22 different sea levels (Section 3.1.1.). This dijit features a mouse-over event that prints a
23 message box of total estimated residential damages. Social vulnerability (Fig. 4b) is output
24 simultaneously with economic vulnerability, and jointly displayed within the "Results" tab of
25 the accordian pane (Fig.2). At a glance, users can assess the socio-economic vulnerabilities of
26 any set of APZ polygons.
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65






Figure 4. Graphical output of economic (a) and social (b) vulnerability data for user-selected planing zone features.

Two other functions provided by the main user interface include a "Search" dijit in the accordian pane, and a "Print" button in the task pane. The search dijit provides a geocoder to support address lookups, while the Print button launches an Adobe compatible portable document format (.pdf) map complete with legend, scale bar, etc.

3.3. Processing Layer

1
2
3
4 Web-SDSS's are not intended to be complete geoanalytical environments, and cannot
5 replace already established geographic information systems (GIS). Nevertheless, vendors
6 such as ESRI have exposed geoprocessing tools as web services, allowing for the execution
7 of complicated and sophisticated geospatial analyses. Such functionality has been
8 incorporated into the CAV, as well as some other web-SDSS, e.g., Rao et al (2007).
9

10
11
12
13 In effect, ArcGIS Server geoprocessing scripts are "stored procedures" which receive
14 data dynamically at runtime. Geoprocessing tasks can be as complicated as need be and,
15 given that they are server-side processes, tend to quite efficient to run. Two geoprocessing
16 tasks were created to support vulnerability analysis within the CAV: EconVuln and
17 SocVuln (Fig. 5). Both of these geoprocessing tasks perform a spatial overlay of user-
18 selected APZ polygons with property parcel data (stored within the Spatial server of the
19 Data Layer, Fig.1), tabulate economic or social vulnerability, and return the output as a result
20 set to be parsed and printed as text and graphical information in the Results window of the
21 Accordian pane (Fig.2).
22
23
24
25
26
27
28
29
30



55 **Figure 5.** Model outlining the functionality of the geoprocessing service SocVuln, which
56 provides a mean summary of the social vulnerability components for land parcels
57 intersecting the user-selected adaptation planning zone(s).
58
59
60
61
62
63
64
65

1
2
3
4 3.4. *Anticipated Workflow During Adaptation Planning Sessions*
5
6
7

- 8 1. Phase 1: unstructured (but not necessarily non-systemic) data exploration.

9
10 Participants expose layers as desired, and inspect regions of the map of greatest
11 concern. There are three dimensions or properties that are likely to effect the
12 discrimination of important patterns or groupings in the data:
13
14

15
16 [visible layers}, map scale, location]
17
18

19
20 Progression may be slow in this phase depending upon the size of the map and the
21 number of data layers to inspect. It is expected that the "information seeking
22 mantra" of Schneiderman (1996), or "overview first, zoom and filter, and then
23 details-on-demand" constitutes a sensible approach for consideration of different
24 data layers, in different locations, at different scales. Users can drill-down and access
25 details as necessary.
26
27
28
29
30

31
32 Other frameworks, for instance, the approach of Flax et al. (2002), could be used to
33 guide the process of hazard identification and analysis. Depending upon the
34 expertise and knowledge of the participant group, there may be a decision to initiate
35 data exploration with the most familiar aspect of community vulnerability, e.g.,
36 physical infrastructure, economic damages, or social vulnerability.
37
38
39
40
41
42

- 43 2. Phase 2: feature creation and annotation. Users identify locations of concern through
44 the feature-creation pane (Fig. 2), and provide "expert" knowledge. Planning zones
45 may delineated at the same time, or as part of a separate process following problem
46 identification of LOCs.
47
48
49
50

- 51
52 3. Phase 3: analysis of social and economic vulnerability of LOCs and assignment of
53 relative priority rankings. Coincident with this is the discussion of mitigation
54 opportunities for lowering risk and increasing resiliency. It is assumed, however, that
55 further analysis of the costs and benefits, as well as social and political feasibility, will
56
57
58
59
60
61
62
63
64
65

1
2
3
4 be conducted outside of the SDSS session. Such an analysis could be conducted, for
5 instance, by independent experts or advisory committees.
6
7
8
9

10 11 12 **4. Adaptation Focus Groups and Software Assessment** 13 14

15
16
17 Six, small-group adaptation focus groups were conducted between January 20 and
18 February 3, 2014. These sessions gathered $n = 31$ expert stakeholders from the Town of
19 Sackville, New Brunswick, and included members of Town council, engineers and
20 emergency measures personnel, town planners, dyke managers, representatives from non-
21 governmental agencies, and community service providers. Small group sessions were
22 adopted in order to: (1) foster knowledge exchange between participants, recognizing that
23 knowledge is, in part, socially constructed (Ramsey 2009); (2) counteract the tendency for
24 self-censorship in larger group settings (see Hopfer and MacEachren 2007); and (3) gather
25 participant feedback on the CAV software.
26
27
28
29
30
31
32

33 To expedite the three-hour sessions, the principal investigator acted as the software
34 “chauffeur” (sensu Aggett and McColl 2006, Jankowski et al. 2006), interacting with the
35 toolkit as directed by participants who used laser pointers to identify locations displayed on a
36 projector screen. This has the advantage of shielding users from the technical details of using
37 the software, and allowed them to focus attention on aspects of community vulnerability.
38 Initially, overviews of all data layers was provided, followed by identification of locations of
39 concern (LOCs), delineation of special adaptation planning zones (APZs), and analysis of
40 the associated economic and social vulnerabilities. Later sessions commenced using the data
41 layers most familiar or interesting to participants, which served as a more effective “ice
42 breaker” and improved the willingness of participants to volunteer information.
43
44
45
46
47
48
49
50

51 Groups identified common as well as unique locations, and discussed different
52 implications for each (Fig. 6). Features included vulnerable sections of dykes and aboiteaux;
53 need for maintenance of drainage ditches; agricultural impacts; commercial and industrial
54 impacts; interruption to community and emergency services; flooded neighbourhoods and
55 neighbourhoods temporarily isolated as a result of road flooding; concerns about land use
56 decisions; questions about the resilience of lift stations and the sewage network; vulnerable
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

historic sites; and, vulnerable populations such as children and elderly. Focus groups identified five main adaptation planning zones, with a total economic exposure of \$6,470,000 (CDN) under the current 10% chance per year, 8.9 metre flood scenario. This estimated damage cost more than doubles to \$13,475,000 (CDN) under a 10 metre flood depth (4% chance per year by 2100), for parcels with a current tax assessed value of about \$21.6 million (CDN).

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 6. Locations of concern (LOCs, filled circles) and adaptation planning zones (APZs, shaded polygons) identified during six adaptation planning sessions (a-e). Also indicated is the possible extent of an 8.9 m flood (CGVD28 vertical datum, hatched area).

1
2
3
4
5
6 A second phase of consultation took the form of a single plenary session on February
7 18, 2014. The chief findings included an assessment of the CAV toolkit (Fig. 7, $n = 28$
8 respondents), new information about flood-risk tolerance, and a set of key recommendations
9 for lowering community vulnerability. Focus group participants reported high levels of
10 satisfaction with aesthetic (Fig. 7b) and navigational aspects of the CAV (Fig. 7d), and
11 evaluated it to be both easy to learn (Fig. 7c) and easy to understand (Fig. 7a). Fewer focus
12 group participants reported the menu to be "friendly" to understand (Fig. 7e), and there were
13 aspects of the flood risk problem that some participants felt were not addressed (Fig. 7f):
14 provisioning of information about inland flooding areas, locations of wetlands, finer
15 topographical information, more information about the sewage collection system, support
16 for photographs, and information about flood depths.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65




Figure 7. Assessment of CAV software by focus group participants ($n = 28$ responses).

1
2
3
4 With regards to flood-risk tolerance, a survey administered at the end of the plenary
5 session ($n = 16$ respondents) indicated that even a moderate 1% chance (1-in-100 year) flood
6 event deters slightly over an estimated 70% of respondents. By the time one considers flood
7 events of 1-in-50 years return frequency (2% chance per year), approximately 90% of
8 respondents indicated a willingness to relocate. This suggests that 1-in-10 year flood maps
9 under represent people's true sensitivity to risk.
10
11
12
13
14
15
16
17

18 **5. Discussion**

19
20
21

22 Given the potential severity of future climate change impacts (IPCC report WGII
23 2014) there is clearly a pressing need for proactive, systematic, community-level assessment
24 of climate change vulnerability (Flax et al. 2002). According to Flax et al. (2002), recognition
25 of the importance of this process led FEMA to tie disaster-relief funding to state-level
26 submission of risk mitigation plans. SDSS software tools can "fast track" the preparation of
27 such plans by aiding visualization, knowledge generation, and analysis of vulnerability, and
28 can work within other pre-existing frameworks. Empirical studies confirm that map-based
29 data reduces the cognitive load on users and leads to faster, more efficient decision making
30 (Crossland et al. 1995, Keenan 2006). In general, the more risk-data available the greater the
31 opportunity to pinpoint high-risk areas (Flax et al. 2002).
32
33
34
35
36
37
38
39

40 SDSS software can facilitate adaptation planning in other ways. As elaborated by Moss
41 et al. (2013), when decision makers confront adaptation challenges, they tend to be bound by
42 considerations of cost, feasibility, social acceptance, tradition, and other factors. This
43 immediately inhibits creative problem solving. Can SDSS help to close the "usability gap" of
44 scientific information and help decision makers whose primary concerns are fixated on other
45 matters? Before answering that question we need to consider the processes an exploratory
46 software tool can support.
47
48
49
50
51
52

53 Koua and Kraak (2004) consider there to be three key exploratory tasks for knowledge
54 construction. The first set of tasks involves categorization and classification, such that
55 attention is drawn to clusters or trends. Perception of these patterns may be facilitated by
56 different perspectives, e.g., 2D and 3D layers, animations. The CAV case study illustrates
57 that interactive visualization of vulnerable infrastructure, with risk layers superimposed on
58
59
60
61
62
63
64
65

1
2
3
4 the same map, is a rapid and efficient way to identify vulnerable locations. In the case study,
5 focus group participants consistently identified some of the vulnerabilities (Fig 6). For
6 example, the sewage lagoon and sections of highway potentially inundated under a 8.9 m
7 flood event were uniformly flagged by participants. Group consensus may naturally emerge
8 under these conditions.
9

10
11
12
13 In addition to supporting user-driven queries about which buildings or features lie
14 within the flood risk zone, users could upload their own knowledge. Through the use of two
15 levels of annotation (LOCs and APZs), users could either "pin" specific locations of concern
16 or classify areas by delineating them using APZ polygons. For instance, participants who
17 were personally familiar with eroded sections of dyke identified those locations on the map,
18 leading to their digitization and, hence, their documentation. Participants benefited from the
19 knowledge exchange, which deepened overall flood-risk awareness of the group.
20
21
22
23
24
25

26 Koua and Kraak's (2004) second exploratory task ("comparison") was facilitated
27 through the social and economic vulnerability output provided by the Processing Layer (Fig.
28 1). Participants gained immediate feedback on which APZ polygons were most at-risk,
29 thereby ensuring that subsequent discussion was informed by a rational appraisal of the
30 evidence.
31
32
33

34 The "reflective" aspect of exploratory visualization (evaluation, integration,
35 generalization) occurs naturally when general statements about vulnerability can be made.
36 For the case study, most locations of concern could either be classified as vulnerable to
37 flooding or vulnerable to isolation.
38
39
40

41 To render SDSS tools effective in collaborative settings, Ramsey (2009) recommends
42 making SDSS tools as flexible as possible. This constitutes an important departure from the
43 knowledge-DSS (see Power and Kaparathi 2002) applications of Best et al. 2007 and Rao et
44 al. 2007, where design goals focused on delivering outputs from complex, custom-designed
45 algorithms applied to well-structured problems. Many aspects of environmental management
46 are qualitative, not quantitative, and involve discursive elements surrounding social and
47 political factors which may, at first blush, appear to be external to the problem domain. Map
48 visualization should be pursued for those aspects of the problem amenable to spatialization;
49 when they cannot, other media (e.g., annotations, text) should be used to capture and
50 represent that knowledge (Risbey 2009). This philosophy was pursued during CAV
51 development, though emphasis was placed on two-dimensional, map-based representation.
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 Previous work (Lieske et al. 2014) identified other forms of visualization (e.g., 3D
5 animations) to be important vehicles for risk communication, but given the high level of
6 problem familiarity of the participants in this project these tools were not deemed worth the
7 time to develop. Nevertheless, depending upon the intended audience, other modes of
8 representation may be effective for communicating particular aspects of the problem domain
9 (though see Lieske 2012 for a caveat regarding the limitations associated with presenting
10 complex visualizations to "casual users").

11
12
13
14
15
16 Jankowski et al. (1997) argue that collaborative problem solving and consensus
17 building stands to benefit from carefully designed decision support systems. To what degree
18 did the CAV contribute to group collaboration and consensus building? The open-ended
19 nature of system, i.e., support for user-contributed information parallel to visualization of
20 quantitative spatial data (along with associated supporting "metadata"), and an easy-to-use
21 and intuitive interface, lent flexibility to the software and allowed it to be applied in new
22 ways. For instance, project participants suggested a wide range of novel applications for the
23 CAV, such as: identification of areas that could be used as flood buffers for water retention,
24 as a teaching tool for schools, or as a means to map vulnerable members of the community
25 (Table 2). Given the fact that SDSS's tend to fail for sociological rather than technological
26 reasons (Rivington et al. 2007), the healthy range of potential applications for the CAV
27 suggests that it possesses a requisite level of sophistication, flexibility and ease-of-use to
28 ensure relevance for a range of potential users.
29
30
31
32
33
34
35
36
37
38
39
40
41

42 Table 2. Potential applications of the CAV SDSS software, as suggested by project
43 participants.
44

-
- 45 ■ [Use] By private citizens to check when Town Council wants to approve a project; the
46 tool could allow citizens to be more aware of Town decisions and enable them to lobby
47 for by-law changes.
 - 48 ■ To identify areas that could be rehabilitated into flood buffers.
 - 49 ■ [Use] By NGOs to look at vulnerable areas and focus work on outreach with vulnerable
50 community members.
 - 51 ■ The tool could be used as a promotional and awareness tool for communities and
52 therefore, as an educational resource.
 - 53 ■ To see where flood boundaries are and what will be affected.
 - 54 ■ For future program planning.
 - 55 ■ For emergency planning, e.g., evacuation or [placement of] reception centre.
 - 56 ■ For land use planning and development.
- 57
58
59
60
61
62
63
64
65

- 1
 - 2
 - 3
 - 4
 - 5
 - 6
 - 7
 - 8
 - 9
 - 10
 - 11
 - 12
 - 13
 - 14
 - 15
 - 16
 - 17
 - 18
 - 19
 - 20
 - 21
 - 22
 - 23
 - 24
 - 25
 - 26
 - 27
 - 28
 - 29
 - 30
 - 31
 - 32
 - 33
 - 34
 - 35
 - 36
 - 37
 - 38
 - 39
 - 40
 - 41
 - 42
 - 43
 - 44
 - 45
 - 46
 - 47
 - 48
 - 49
 - 50
 - 51
 - 52
 - 53
 - 54
 - 55
 - 56
 - 57
 - 58
 - 59
 - 60
 - 61
 - 62
 - 63
 - 64
 - 65
-
- To lead discussions and to stimulate conversation.
 - To help when buying or selling land or houses.
 - To help make elected officials aware of potential flooding.
 - To help municipal councils and planning commissions in creating adaptation plans.
 - When constructing infrastructure and determining where to invest in infrastructure repairs or upgrades.
 - To evaluate roadway systems.
 - To evaluate how many homes and businesses may be affected during an event.
 - For public presentations and meetings.
 - For calculating risk.
 - To introduce flood issues to new communities beyond Sackville.
 - As a teaching tool
 - For advance warnings during major weather events, such as storm surges, rainfall, high winds, and high tides.
 - To plot people in the community that use home care.
 - To understand the risks involved with flooding in a particular area of town or on [Mount Allison University] campus.

An important component of adaptation planning involves priority setting and scheduling. While the CAV supported simple priority ranking, e.g., attribution of an integer rank to adaptation planning zones (Fig.3b), participants in the case study were hesitant to volunteer values. This suggests that a different prioritization process was necessary to support this activity. Provision of an "analytical hierarchical process" (e.g., Jankowski et al. 1997, Karnatak et al. 2007) may facilitate prioritization, with individual APZs being compared in a pair-wise manner to help assign relative importance. An anonymous voting system may also have encouraged participation. However, given the complexity of multi-stakeholder, multi-dimensional problems such as community vulnerability to climate change, there may be no simple approach to setting priorities (Ramsey 2009). For instance, how do we resolve the social implications of ranking one neighbourhood in "greater" need of risk reduction intervention than another, with the attendant (and implicit) value system influencing the appraisal process? Such issues are not resolvable purely through the use of SDSS software.

Successful identification and prioritization of candidate adaptation measures may not, in itself, lead to implementation. Individual communities may not have access to the resources necessary to carry out significant adaptation measures, i.e., those with greatest chance of lowering community vulnerability. In the case study, relocation of the core of vulnerable

1
2
3
4 residences via land acquisition was estimated to exceed \$21 million (CDN). Participants
5 concluded that even if such an initiative could be enacted, other system components would
6 remain vulnerable, e.g., the travel corridor comprised of the Trans-Canada Highway and CN
7 railway, and local agricultural activity. Recognition of these linkages led to the conclusion
8 that the existing dyke system, with a relatively modest investment of approximately \$2.5
9 million (CDN), was an efficient option for counter-acting sea level rise in the short term and
10 protecting the range of community assets. Over the long-term (approximately 40 years),
11 limitations in the capacity to raise the height of the dykes without major re-engineering
12 would likely nullify their capacity to withstand future sea levels. As a consequence of these
13 findings, participants quickly identified pro-active emergency response and land use planning
14 as vital to limiting and lowering community vulnerability in the interim.

15
16 While the case study involved expert stakeholders, i.e., those whose professional
17 responsibilities or areas of knowledge directly pertained to coastal flood risk reduction, the
18 CAV could just as easily been used as an outreach and information gathering tool and
19 addressed a wider cross-section of the general public. In fact, it has been argued that
20 fostering multi-stakeholder, iterative, consultative discussions are a good remedy for
21 preventing decision making processes becoming unmanageable or contentious (Jankowski et
22 al. 2006, Nyerges et al. 2006). GIS and the Web hold great potential for public use (Kingston
23 et al. 2000). Rather than attempt to force top-down, "winner vs. loser" models, SDSS's have
24 the potential to involve everyone in risk appraisal visualization, possibly resulting in more
25 equitable, consensus-based sets of recommendations endorsed by most or all stakeholders
26 (Flax et al. 2002, Nyerges et al. 2006).

27
28 While the case study reported here was for a small municipality, the CAV software can
29 be applied at any scale, for any hazard for which spatial information is available. The CAV
30 was demonstrated to be a rich environment for spatial analysis, allowing submission of
31 geographically-specific knowledge as well as rapid assessment of social and economic
32 vulnerability at particular locations. Such a process supports objective and rational risk
33 appraisal and stands to better inform stakeholders. Given the pressing nature of climate
34 change, and the long time frame required to fulfill some adaptation plans, it is crucial that
35 communities utilize all tools at their disposal to immediately develop and initiate adaptation
36 strategies.

Acknowledgements

The author acknowledges the financial support of the New Brunswick Environmental Trust Fund (ETF) and the Social Sciences and Humanities Research Council (SSHRC) Small Universities Grant. The author acknowledges the data management provided by J. Bornemann, and facilitation of community focus groups by L.A. Roness and E.A. Phillips. The author is grateful to A. Hamilton-Wright for kindly reviewing an earlier draft of the manuscript. The author also thanks the participants who generously donated their time to evaluate the CAV, while also providing important flood adaptation recommendations for the Sackville community. Lastly, the author acknowledges the many software developers who have contributed the free and open-source software libraries used in this project.

Figure Legends

Figure 1. System architecture of the Community Adaptation Viewer (CAV).

Figure 2. Schema of the main user interface.

Figure 3. Text controls for editing attribute information for user-created locations of concern (a), and adaptation planning zones (b).

Figure 4. Graphical output of economic (a) and social (b) vulnerability data for user-selected planing zone features.

Figure 5. Model outlining the functionality of the geoprocessing service SocVuln, which provides a mean summary of the social vulnerability components for land parcels intersecting the user-selected adaptation planning zone(s).

Figure 6. Locations of concern (LOCs, filled circles) and adaptation planning zones (APZs, shaded polygons) identified during six adaptation planning sessions (a-e). Also indicated is the possible extent of an 8.9 m flood (CGVD28 vertical datum, hatched area).

Figure 7. Assessment of CAV software by focus group participants ($n = 28$ responses).

REFERENCES

- Aggett, G., C. McColl, 2006. Evaluating decision support systems for PPGIS applications. *Cartography and Geographic Information Science* 33: 77-92.
- Andrienko, N. and G. Andrienko. 2006. *Exploratory analysis of spatial and temporal data: a systematic approach*. Springer, New York.
- Andrienko, G., Andrienko, N., Jankowski, P., Keim, D., Kraak, M.-J., MacEachren, A., and S. Wrobel. 2007. Geovisual analytics for spatial decision support: Setting the research agenda. *International Journal of Geographical Information Science* 21: 839–857.
- Bhargava, H.K., D.J. Power and D. Sun. 2007. Progress in Web-based decision support technologies. *Decision Support Systems* 43: 1083-1095.
- Best, B.D., P.N. Halpin, E. Fukioka, A.J. Read, S.S. Qian, L.J. Hazen and R.S. Schick. 2007. Geospatial web services within a scientific workflow: predicting marine mammal habitats in a dynamic environment. *Ecological Informatics* 2: 210-223.
- Chen, C. 2005. Top ten unsolved information visualization problems. *IEEE Computer Graphics and Applications* July/August: 12-16.
- Chen, M., Y. Liou, C.-W. Wang, Y.-W. Fan, Y.-P. J. Chi. 2007. TeamSpirit: design, implementation, and evaluation of a web-based group decision support system. *Decision Support Systems* 43: 1186-1202.
- Crossland, M.D., B.E. Wynne and W.C. Perkins. 1995. Spatial decision support systems: An overview of technology and a test of efficacy. *Decision Support Systems* 14: 219-235.
- Cutter, S.L., J.T. Mitchell, and M.S. Scott. 2000. Revealing the vulnerability of people and places: a case study of Georgetown County, South Carolina. *Annals of the Association of American Geographers* 90: 713-737.
- Eppler, M.J. and M. Aeschmann. 2009. A systematic framework for risk visualization in risk management and communication. *Risk Management* 11: 67-89.
- ESRI, 2014. ArcGIS Server, <http://www.esri.com/software/arcgis/arcgisserver> (Downloaded May 2014).
- Flanagan, D. 2011. *Javascript: The definitive guide*. O'Reilly, Sebastopol, California.
- Flax, L., R. Jackson and D. Stein. 2002. Community vulnerability assessment tool methodology. *Natural Hazards Review* 3: 163-176.
- Füssel, H-M. 2007. Vulnerability: a generally applicable conceptual framework for climate change research. *Global Environmental Change* 17: 155-167.
- Gulliksen, J., G. Bengt, I. Boivie, S. Blomkvist, J., Perrson and Å. Cajander. 2003. Key principles for user-centred systems design. *Behaviour & Information Technology* 22: 397-409.
- Hebb, A. and L. Mortsch. 2007. Floods: mapping vulnerability in the Upper Thames watershed under a changing climate. http://zerofootprint.uwaterloo.ca/research/aird/aird_pub/Upper_Thames_Vulnerability_Mapping_Final_Report.pdf (Downloaded Jan 16, 2014).
- Hopfer, S. and A.M. MacEachren. 2007. Leveraging the potential of geospatial annotations for collaboration: a communication theory perspective. *International Journal of Geographical Information Science* 21: 921-934.
- IPCC WGII AR5, 2014. *Climate change 2014: impacts, adaptation, and vulnerability. Summary for Policymaker*, <http://ipcc->

- 1
2
3
4 wg2.gov/AR5/images/uploads/IPCC_WG2AR5_SPM_Approved.pdf (Downloaded
5 May, 2014).
- 6
7 Jankowski, P., T.L. Nyerges, A. Smith, T.J. Morre, and E. Horvath. 1997. Spatial group
8 choice: a tool for collaborative spatial decision-making. *International Journal of*
9 *Geographical Information Science* 11: 577-602.
- 10 Jankowski, P., T. Nyerges, S. Robischon, K. Ramsey, and D. Tuthill. 2006. Design
11 considerations and evaluation of a collaborative, spatio-temporal decision support
12 system. *Transactions in GIS* 10: 335-354.
- 13
14 Karnatak, H.C., S. Saran, K. Bhatia, and P.S. Roy. 2007. Multicriteria spatial decision analysis
15 in web GIS environment. *Geoinformatica* 11: 407-429.
- 16
17 Keenan, P.B. 2006. Spatial decision support systems: a coming of age. *Control and*
18 *Cybernetics* 35: No.1.
- 19
20 King, D., and C. MacGregor. 2000. Using social indicators to measure community
21 vulnerability to natural hazards. *Australian Journal of Emergency Management* 15(3): 52-
22 57.
- 23
24 Kingston, R., S. Carver, A. Evans and I. Turton. 2000. Web-based public participation
25 geographical information systems: an aid to local environmental decision-making.
26 *Computers, Environment and Urban Systems* 24: 109-125.
- 27
28 Koua, E.L. and M-J. Kraak. 2006. A useability framework for the design and evaluation of
29 an exploratory geovisualization environment. *Proceedings of the Eighth International*
30 *Conference on Information Visualization (IV'04)*,
31 http://www.geovista.psu.edu/publications/2004/IEEEpaperUsability_Koua.pdf
32 (Downloaded May, 2014).
- 33
34 Lieske, D.J. 2012. Towards a framework for designing spatial and non-spatial visualizations
35 for communicating climate change risks. *Geomatica* 66: 255-265.
- 36
37 Lieske, D.J., T. Wade and L.A. Roness. 2014. Climate change awareness and strategies for
38 communicating the risk of coastal flooding: a Canadian Maritime case example.
39 *Estuarine and Coastal Shelf Science* 140: 83-94.
- 40
41 Maguire, M. 2001. Methods to support human-centred design. *International Journal of*
42 *Human-Computer Studies* 55: 587-634.
- 43
44 Mennis, J. and D.J. Peuquet. 2003. The role of knowledge representation in geographic
45 knowledge discovery: a case study. *Transactions in GIS* 7: 371-391.
- 46
47 Moss et al. 2013. Hell and high water: practice-relevant adaptation science. *Science* 342: 696-
48 698.
- 49
50 National Research Council. 2006. *Facing hazards and disasters: understanding the human*
51 *dimensions*. National Academies Press, Washington, D.C.
- 52
53 Nyerges, T., P. Jankowski, D. Tuthill, and K. Ramsey. 2006. Collaborative water resource
54 decision support: results of a field experiment. *Annals of the American Association of*
55 *Geographers* 96: 699-725.
- 56
57 Peuquet, D.J. and M-J. Kraak. 2002. Geobrowsing: creative thinking and knowledge
58 discovery using geographic visualization. *Information Visualization* 1: 80-91.
- 59
60 Power, D.J. and S. Kaparathi. 2002. Building web-based decision support systems. *Studies in*
61 *Informatics and Control* 11: 291-302.
- 62
63 Ramsey, K. 2009. GIS, modeling, and politics: on the tensions of collaborative decision
64 support. *Journal of Environmental Management* 90: 1972-1980.
- 65
66 Rao, M., G. Fan, J. Thomas, G. Cherian, V. Chudiwale, M. Awawdeh. 2007. A web-based
67 GIS decision support system for managing and planning USDA's Conservation
68 Reserve Program (CRP). *Environmental Modelling & Software* 22: 1270-1280.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Rinner, C. 2003. Web-based spatial decision support: status and research directions. *Journal of Geographic Information and Decision Analysis* 7: 14-31.

Risbey, J., M. Kandlikar, H. Dowlatabadi, and D. Graetz. 1999. Scale, context, and decision making in agricultural adaptation to climate variability and change. *Mitigation and Adaptation Strategies for Global Change* 4: 137-165.

Rivington, M., K.B. Matthews, G. Bellocchi, K. Buchan, C.O. Stöckle and M. Donatelli. 2007. An integrated assessment approach to conduct analyses of climate change impacts on whole-farm systems. *Environmental Modelling & Software* 27: 202-210.

Roness, L.A. 2013. New Brunswick climate change communication needs assessment: expanding the Tantramar Map Viewer. <http://arcgis.mta.ca/docs/Tantramar%20Viewer%20Project%20-%20Report%20-%20FINAL.pdf> [downloaded 24 April 2014].

Russell, M.A. 2008. *Dojo: The definitive guide*. O'Reilly, Sebastopol, California.

Schniederman, B. 1996. The eyes have it: a task by data type taxonomy for information visualization. *Proceedings of Visual Languages* 96.

Sugumaran, R., and J. DeGroot. 2011. *Spatial decision support systems: principles and practices*. CRC Press.

Wilby, R.L. and R. Keenan. 2012. Adapting to flood risk under climate change. *Progress in Physical Geography* 36: 348-378.

Wilson, J., R. Trenholm, J. Bornemann and D. Lieske. 2012. Forecasting economic damages from storm surge flooding: a case study in the Tantramar region of New Brunswick <http://arcgis.mta.ca/docs/ForecastingEconomicDamagesfromTantramarStormSurgeFlooding.pdf> (Downloaded May, 2014).

Wu, S-Y., B. Yarnal and A. Fisher. 2002. Vulnerability of coastal communities to sea-level rise: a case study of Cape May County, New Jersey, USA. *Climate Research* 22: 255-270.